


Rations for Chickens *

For small scale commercial chickens, fed full-time, one of the most important aspects we need to know is the crude protein (CP) level of the ration. Generally, we start with a high level of protein and later on, as the bird gets older, we can reduce the protein level. Many small scale chicken producers in Malawi follow the following general guidelines for CP content of rations:

| Broilers | Layers | CP |
|----------------------------|---------------------------|--------|
| Starter ration, 0-2 weeks | Starter ration, 0-6 weeks | 20-22% |
| Grower ration, 2-4 weeks | Grower ration, 6-20 weeks | 18-20% |
| Finisher ration, 4 weeks + | Layer ration, 20 weeks + | 16-18% |

Don't change rations suddenly. There is usually no need. The following diagram shows how to gradually change from one ration to another over a few days.



We can calculate the contribution of various feed components to the nutrient content of the ration. The approximate nutrient values of many feed components can be obtained from published data (such as is available through the joint project "Feedipedia"). Using this information, we can design home-mixed rations using locally available ingredients. This may enable a significant cost saving over commercially prepared complete rations. The first step in this procedure is to determine what ingredients are available locally.

The following table shows a few sample rations which may be suitable for supplementary feeding in Malawi and the approximate estimated crude protein content of the respective ration. The ingredients in these rations are available in Malawi at least in some localities. Other nutritional information about these rations such as energy content is not shown here. These calculations are based on the estimated CP and moisture content of the ingredients, which may vary from supplier to supplier. It is therefore necessary to take care in using these rations. A proper nutritional analysis of the prepared ration is essential if significant numbers or valuable poultry are involved.

On request, RPC can tailor rations like these to ensure the least cost mix of ingredients. This process is called "least cost ration analysis".



* Copy, distribute, but please attribute. This document 'Rations for chickens' by the Rural Poultry Centre in Malawi is licensed under a [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).

Examples of Rations

| | Ration 1 | Ration 2 | Ration 3 | Ration 4 | Ration 5 | Ration 6 |
|--------------------------------|-----------|-----------|-----------|-----------|-----------|-----------|
| Ingredient | Kg/100 kg | Kg/100 kg | Kg/100 kg | Kg/100 kg | Kg/100 kg | Kg/100 kg |
| Madeya | 50 | 42 | 43.6 | 40 | 37 | 30.8 |
| Crushed maize | 37.6 | 43.5 | 38 | 40.7 | 39 | 34 |
| Fish meal/dried fish | 10.5 | - | 6 | - | 1.5 | - |
| Soybean meal | - | 11.2 | 10 | 11 | 20 | 20 |
| Rice bran | - | - | - | - | - | 7 |
| Groundnut meal (afatoxin free) | - | - | - | 5 | - | 5 |
| MCP | 0.3 | 0.7 | 0.4 | 0.6 | 2.0 | 0.1 |
| Lime | 1.0 | 2.0 | 1.4 | 2.1 | 0.4 | 2.5 |
| Salt | 0.3 | 0.3 | 0.3 | 0.3 | 0.3 | 0.3 |
| Premix | 0.3 | 0.3 | 0.3 | 0.3 | 0.3 | 0.3 |
| Total | 100 | 100 | 100 | 100 | 100 | 100 |
| Est. CP | 14% | 14% | 16% | 16% | 18% | 20.0% |

How much will my chickens eat?

It is difficult to be precise about how much total food chickens will eat. There are many factors which affect the total quantity of feed consumed including:

- Feed factors: quality of the feed;
- Bird factors: sex, age, breed, and type of chicken, as well as health status;
- Environmental factors: temperature conditions, stress levels, management.

Subject to variations due to these and similar factors, some published figures for feed consumption are as follows:

Typical Body Weight and Feed Requirements of Broiler Chickens

| Age (wks) | Body wt (grams) | Weekly feed (g) | Cumulative feed (g) |
|-----------|-----------------|-----------------|---------------------|
| 1 | 150 | 132 | 132 |
| 2 | 358 | 281 | 413 |
| 3 | 653 | 467 | 880 |
| 4 | 1,025 | 671 | 1,551 |
| 5 | 1,461 | 848 | 2,404 |
| 6 | 1,914 | 1,070 | 3,475 |
| 7 | 2,363 | 1,179 | 4,654 |
| 8 | 2,790 | 1,297 | 5,951 |
| 9 | 3,198 | 1,411 | 7,362 |

Typical Body Weight and Feed Requirements of Layer Pullets (brown eggs)

| Age (wks) | Body wt (grams) | Weekly feed (g) | Cumulative feed (g) |
|-----------|-----------------|-----------------|---------------------|
| 0 | 36 | 136 | 136 |
| 2 | 118 | 318 | 454 |
| 4 | 327 | 562 | 1,016 |
| 6 | 499 | 699 | 1,715 |
| 8 | 748 | 762 | 2,477 |
| 10 | 898 | 798 | 3,275 |
| 12 | 1,102 | 844 | 4,119 |
| 14 | 1,238 | 898 | 5,017 |
| 16 | 1,379 | 943 | 5,960 |
| 18 | 1,501 | 998 | 6,958 |
| 20 | 1,601 | 1,098 | 8,056 |